


DIABETES IN WOMEN

A heavy toll across the life course



4.5 years

later diagnosis than men for diabetes and other metabolic diseases⁹



A

13%

greater risk of all-cause mortality compared to men living with diabetes⁶



Women with abnormal HbA1c values were




3x

more likely to develop yeast infections than women with normal values⁷

Women living with Type 1 Diabetes are


4x

more likely to develop pre-eclampsia than women without diabetes³



1 in 7

live births in Europe are affected by hyperglycaemia in pregnancy⁴



1/2

of women with GDM* will develop Type 2 Diabetes (T2D) within 5 years of giving birth.¹

Their children are **6x** more likely to develop T2D and childhood obesity compared to children born to mothers without GDM¹

*Gestational Diabetes Mellitus

A 30%

greater excess risk of CVD mortality than men living with diabetes⁶



2x

more likely to develop eating disorders than if they didn't live with diabetes⁵



Women living with diabetes have a **1/3**

higher likelihood of developing depression compared to men living with diabetes²



Women living with T2D have a

19%

greater risk to develop vascular dementia than men living with T2D⁸

● Non-European/Global Studies

Sources

1. Women and Diabetes in the EU; European Institute of Women's Health (EIWH); Policy Brief Eurohealth
2. Sartorius, N. (2018). Depression and diabetes. *Dialogues in Clinical Neuroscience*, 20(1), 47–52. <https://doi.org/10.31887/dcn.2018.20.1/sartorius>
3. Callesen, N., Vestgaard, M., Nørgaard, S. K., Damm, P., Mathiesen, E. R., & Ringholm, L. (2023). Prediction and prevention of preeclampsia in women with preexisting diabetes: the role of home blood pressure, physical activity, and aspirin. *Frontiers in Endocrinology*, 14. <https://doi.org/10.3389/fendo.2023.1166884>
4. MEPs Mobilising for Diabetes; IDF Atlas factsheet – Diabetes in Europe 2021
5. Gagnon, C., Aimé, A., Bélanger, C. Predictors of Comorbid Eating Disorders and Diabetes in People with Type 1 and Type 2 Diabetes. *Can J Diabetes*, 2017 Feb;41(1):52-57. doi: 10.1016/j.jcjd.2016.06.005. Epub 2016 Sep 7. PMID: 27614805.
6. Wang, Y., O'Neil, A., Jiao, Y. et al. Sex differences in the association between diabetes and risk of cardiovascular disease, cancer, and all-cause and cause-specific mortality: a systematic review and meta-analysis of 5,162,654 participants. *BMC Med* 17, 136 (2019). <https://doi.org/10.1186/s12916-019-1355-0>
7. De Leon, E. M., Jacober, S. J., Sobel, J. D., & Foxman, B. (2002). Prevalence and risk factors for vaginal Candida colonization in women with type 1 and type 2 diabetes. *BMC Infectious Diseases*, 2(1). <https://doi.org/10.1186/1471-2334-2-1>
8. Chatterjee, S., Peters, S.A., Woodward, M., Mejia Arango, S., Batty, G.D., Beckett, N., Beiser, A., Borenstein, A.R., Crane, P.K., Haan, M., Hassing, L.B., Hayden, K.M., Kiyohara, Y., Larson, E.B., Li, C.Y., Ninomiya, T., Ohara, T., Peters, R., Russ, T.C., Seshadri, S., Strand, B.H., Walker, R., Xu, W., Huxley, R.R. Type 2 Diabetes as a Risk Factor for Dementia in Women Compared With Men: A Pooled Analysis of 2.3 Million People Comprising More Than 100,000 Cases of Dementia. *Diabetes Care*, 2016 Feb;39(2):300-7. doi: 10.2337/dc15-1588. Epub 2015 Dec 17. PMID: 26681727; PMCID: PMC4722942.
9. Westergaard, D., Moseley, P. L., Serup, F. K. H., Baldi, P., & Brunak, S. (2019). Population-wide analysis of differences in disease progression patterns in men and women. *Nature Communications*, 10(1). <https://doi.org/10.1038/s41467-019-08475-9>